



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/255, 31/35, 31/7048, A61P 3/10		A1	(11) International Publication Number: WO 00/61139
			(43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/08404 (22) International Filing Date: 20 March 2000 (20.03.00) (30) Priority Data: 60/128,398 8 April 1999 (08.04.99) US (71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route #202, P.O. Box 300, Raritan, NJ 08869-0602 (US). (72) Inventor: COTTRELL, Sandra, C.; 2 Granite Circle, Doylestown, PA 18901 (US). (74) Agents: CIAMPORCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: ANTICONVULSANT DERIVATIVES USEFUL IN REDUCING BLOOD GLUCOSE LEVELS			
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>(I)</p> </div> <div style="text-align: center;"> <p>(II)</p> </div> </div>			
(57) Abstract Use of anticonvulsant derivatives of formula (I) for reducing blood glucose levels, wherein X is CH or oxygen; R ₁ is hydrogen or alkyl; and R ₂ , R ₃ , R ₄ and R ₅ are independently hydrogen or alkyl and, when X is CH ₂ , R ₄ and R ₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R ₂ and R ₃ and/or R ₄ and R ₅ together may be a methylenedioxy group of formula (II), wherein R ₆ and R ₇ are the same or different and are hydrogen, or alkyl and are joined to form a cyclopentyl or cyclohexyl ring.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ANTICONVULSANT DERIVATIVES USEFUL IN REDUCING BLOOD GLUCOSE LEVELS

BACKGROUND OF THE INVENTION

5

Compounds of Formula I:



are structurally novel antiepileptic compounds that are highly effective anticonvulsants
10 in animal tests (Maryanoff, B.E, Nortey, S.O., Gardocki, J.F., Shank, R.P. and
Dodgson, S.P. *J. Med. Chem.* 30, 880-887, 1987; Maryanoff, B.E., Costanzo, M.J.,
Shank, R.P., Schupsky, J.J., Ortegon, M.E., and Vaught J.L. *Bioorganic & Medicinal
Chemistry Letters* 3, 2653-2656, 1993). These compounds are covered by US Patent
No.4,513,006. One of these compounds 2,3:4,5-bis-O-(1-methylethylidene)- β -D-
15 fructopyranose sulfamate known as topiramate has been demonstrated in clinical trials
of human epilepsy to be effective as adjunctive therapy or as monotherapy in treating
simple and complex partial seizures and secondarily generalized seizures (E. FAUGHT,
B.J. WILDER, R.E. RAMSEY, R.A. REIFE, L D. KRAMER, G.W. PLEDGER, R.M.
KARIM et. al., *Epilepsia* 36 (S4) 33, 1995; S.K. SACHDEO, R.C. SACHDEO, R.A.
20 REIFE, P. LIM and G. PLEDGER, *Epilepsia* 36 (S4) 33, 1995), and is currently
marketed for the treatment of simple and complex partial seizure epilepsy with or
without secondary generalized seizures in approximately twenty countries including the
United States, and applications for regulatory approval are presently pending in several
additional countries throughout the world.


25 Compounds of Formula I were initially found to possess anticonvulsant activity
in the traditional maximal electroshock seizure (MES) test in mice (SHANK, R.P.,
GARDOCKI, J.F., VAUGHT, J.L., DAVIS, C.B., SCHUPSKY, J.J., RAFFA, R.B.,
DODGSON, S.J., NORTEY, S.O., and MARYANOFF, B.E., *Epilepsia* 35 450-460,
1994). Subsequent studies revealed that Compounds of Formula I were also highly

effective in the MES. test in rats. More recently topiramate was found to effectively block seizures in several rodent models of epilepsy (J. NAKAMURA, S. TAMURA, T. KANDA, A. ISHII, K. ISHIHARA, T. SERIKAWA, J. YAMADA, and M. SASA, Eur. J. Pharmacol. 254 83-89, 1994), and in an animal model of kindled epilepsy (A. WAUQUIER and S. ZHOU, Epilepsy Res. 24, 73-77, 1996).

Clinical studies on topiramate have revealed previously unrecognized pharmacological properties which suggest that topiramate will be effective in the reduction of blood glucose in animals, including but not limited to humans.

10 DISCLOSURE OF THE INVENTION

Accordingly, it has been found that compounds of the following formula I:




15

wherein X is O or CH₂, and R₁, R₂, R₃, R₄ and R₅ are as defined hereinafter are useful in maintaining weight loss.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIEMENTS

20

The sulfamates of the invention are of the following formula (I):




wherein

25 X is CH₂ or oxygen;

R₁ is hydrogen or alkyl; and

R₂, R₃, R₄ and R₅ are independently hydrogen or alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃ and/or R₄ and R₅ together may be a methylenedioxy group of the following

5 formula (II):



wherein

R₆ and R₇ are the same or different and are hydrogen, lower alkyl or are alkyl and are
10 joined to form a cyclopentyl or cyclohexyl ring.


R₁ in particular is hydrogen or alkyl of about 1 to 4 carbons, such as methyl, ethyl and iso-propyl. Alkyl throughout this specification includes straight and branched chain alkyl. Alkyl groups for R₂, R₃, R₄, R₅, R₆ and R₇ are of about 1 to 3 carbons and include methyl, ethyl, iso-propyl and n-propyl. When X is CH₂, R₄ and R₅ may
15 combine to form a benzene ring fused to the 6-membered X-containing ring, i.e., R₄ and R₅ are defined by the alkatrienyl group =C-CH=CH-CH=.

A particular group of compounds of formula (I) is that wherein X is oxygen and both R₂ and R₃ and R₄ and R₅ together are methylenedioxy groups of the formula (II), wherein R₆ and R₇ are both hydrogen both alkyl or combine to form a spiro
20 cyclopentyl or cyclohexyl ring, in particular where R₆ and R₇ are both alkyl such as methyl. A second group of compounds is that wherein X is CH₂ and R₄ and R₅ are joined to form a benzene ring. A third group of compounds of formula (I) is that wherein both R₂ and R₃ are hydrogen.

The compounds of formula (I) may be synthesized by the following methods:

25 (a) Reaction of an alcohol of the formula RCH₂OH with a chlorosulfamate of the formula ClSO₂NH₂ or ClSO₂NHR₁ in the presence of a base such as potassium *n*-butoxide or sodium hydride at a temperature of about -20° to 25° C and in a solvent

such as toluene, THF or dimethylformamide wherein R is a moiety of the following formula (III):



(b) Reaction of an alcohol of the formula RCH_2OH with sulfurylchloride of the formula SO_2Cl_2 in the presence of a base such as triethylamine or pyridine at a temperature of about -40° to 25° C in a solvent such as diethyl ether or methylene chloride to produce a chlorosulfate of the formula RCH_2OSO_2Cl .

The chlorosulfate of the formula RCH_2OSO_2Cl may then be reacted with an amine of the formula R_1NH_2 at a temperature of about 40° to 25° C in a solvent such as methylene chloride or acetonitrile to produce a compound of formula (I). The reaction conditions for (b) are also described by T. Tsuchiya et al. in Tet. Letters, No. 36, p. 3365 to 3368 (1978).

(c) Reaction of the chlorosulfate RCH_2OSO_2Cl with a metal azide such as sodium azide in a solvent such as methylene chloride or acetonitrile yields an azidosulfate of the formula $RCH_2OSO_2N_3$ as described by M. Hedayatullah in Tet. Lett. p. 2455-2458 (1975). The azidosulfate is then reduced to a compound of formula (I) wherein R_1 is hydrogen by catalytic hydrogenation, e.g. with a noble metal and H_2 or by heating with copper metal in a solvent such as methanol.

The starting materials of the formula RCH_2OH may be obtained commercially or as known in the art. For example, starting materials of the formula RCH_2OH wherein both R_2 and R_3 and R_4 and R_5 are identical and are of the formula (II) may be obtained by the method of R. F. Brady in Carbohydrate Research, Vol. 14, p. 35 to 40 (1970) or by reaction of the trimethylsilyl enol ether of a R_6COR_7 ketone or aldehyde with fructose at a temperature of about 25° C, in a solvent such a halocarbon, e.g. methylene chloride in the presence of a protic acid such as hydrochloric acid or a Lewis Acid such as zinc chloride. The trimethylsilyl enol ether reaction is described by G. L. Larson et al in J. Org. Chem. Volaa 38, No. 22, p. 3935 (1973).

Further, carboxylic acids and aldehydes of the formulae $RCOOH$ and $RCHO$ may be reduced to compounds of the formula RCH_2OH by standard reduction techniques, e.g. reaction with lithium aluminum hydride, sodium borohydride or

borane-THF complex in an inert solvent such a diglyme, THF or toluene at a temperature of about 0° to 100° C, e.g. as described by H.O. House in "Modern Synthetic Reactions", 2nd Ed., pages 45 to 144 (1972).

5 The compounds of formula I: may also be made by the process disclosed in 5,387,700, which is incorporated by reference herein.

The compounds of formula I include the various individual isomers as well as the racemates thereof, e.g., the various alpha and beta attachments, i.e., below and above the plane of the drawing, of R₂, R₃, R₄ and R₅ on the 6-membered ring. Preferably, the oxygene of the methylenedioxy group (II) are attached on the same side
10 of the 6-membered ring.

In a retrospective analysis of the long term open-label studies in epilepsy in 1319 patients, treated with a mean daily dosage of 622 mg/day, for a mean duration of 689 days, 14 subjects were found to be overweight (defined as weight >100 kg) and with high blood glucose levels (>115 mg/dl). These patients had a 5.8% reduction in
15 blood glucose during the period of treatment with topiramate.

For reducing glucose levels in the blood of mammals, a compound of formula (I) may be employed at a daily dose in the range of about 100 mg to 400 mg, usually in two daily divided doses, for an average adult human. A unit dose would contain about 15 to 200 mg of the active ingredient


20 To prepare the pharmaceutical compositions of this invention, one or more sulfamate compounds of formula (I) are intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral, by suppository, or parenteral. In preparing the
25 compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules and tablets, suitable carriers and additives
30 include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated

or enteric coated by standard techniques. Suppositories may be prepared, in which case cocoa butter could be used as the carrier. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable solutions may also be prepared in which case appropriate stabilizing agents may be employed. Topiramate is currently available for oral administration in round tablets containing 25 mg, 100 mg or 200 mg of active agent. The tablets contain the following inactive ingredients: lactose hydrous, pregelatinized starch, microcrystalline cellulose, sodium starch glycolate, magnesium stearate, purified water, carnauba wax, hydroxypropyl methylcellulose, titanium dioxide, polyethylene glycol, synthetic iron oxide, and polysorbate 80.

The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder injection, teaspoonful, suppository and the like from about 25 to about 200 mg of the active ingredient.

WHAT IS CLAIMED IS:

1. A method for reducing blood glucose levels in a mammal comprising administering
 5 to such a mammal a therapeutically effective amount for treating such condition of a compound of the formula I:




wherein

X is CH₂ or oxygen;

- 10 R₁ is hydrogen or alkyl; and

R₂, R₃, R₄ and R₅ are independently hydrogen or alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃ and/or R₄ and R₅ together may be a methylenedioxy group of the following formula (II):

15



wherein

- 20 R₆ and R₇ are the same or different and are hydrogen, or alkyl and are joined to form a cyclopentyl or cyclohexyl ring.

2. The method of claim 1 wherein the compound of formula I is topiramate.
 3. The method of claim 1, wherein the therapeutically effective amount is of from about 100 to 400 mg.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 00/08404

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/255 A61K31/35 A61K31/7048 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, MEDLINE, BIOSIS, EMBASE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 44581 A (GIBBS IRWIN S ;KOTWAL PRAMOD M (US); THAKUR MADHAV S (US); ORTHO M) 10 September 1999 (1999-09-10) abstract page 5, line 19 - line 24 ---	1-3
A	WO 98 00130 A (ORTHO PHARMA CORP) 8 January 1998 (1998-01-08) the whole document ---	1-3
A	US 4 513 006 A (MARYANOFF BRUCE E ET AL) 23 April 1985 (1985-04-23) cited in the application the whole document --- -/--	1-3

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 August 2000

Date of mailing of the international search report

22/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P

INTERNATIONAL SEARCH REPORT

Int ernational Application No

PCT/US 00/08404

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>KYOWA HAKKO: "TOPIRAMATE" DRUGS OF THE FUTURE, ES, BARCELONA, vol. 21, no. 4, 1996, pages 463-465, XP002043895 ISSN: 0377-8282 the whole document -----</p>	1-3

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/08404

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9944581 A	10-09-1999	AU 2798999 A	20-09-1999
WO 9800130 A	08-01-1998	AU 3957897 A	21-01-1998
		CA 2258893 A	08-01-1998
		CZ 9804278 A	11-08-1999
		EP 0915697 A	19-05-1999
		NO 986052 A	23-02-1999
		ZA 9705772 A	28-12-1998
US 4513006 A	23-04-1985	AT 36149 T	15-08-1988
		AU 564842 B	27-08-1987
		AU 3350484 A	04-04-1985
		CA 1241951 A	13-09-1988
		DE 3473143 D	08-09-1988
		DK 198191 A,B,	09-12-1991
		DK 198291 A	09-12-1991
		DK 457784 A,B,	27-03-1985
		EP 0138441 A	24-04-1985
		ES 536225 D	16-11-1985
		ES 8602634 A	16-03-1986
		FI 843765 A,B,	27-03-1985
		HU 36784 A,B	28-10-1985
		IE 57684 B	24-02-1993
		JP 1804249 C	26-11-1993
		JP 5005824 B	25-01-1993
		JP 60109558 A	15-06-1985
		JP 5331132 A	14-12-1993
		KR 9201775 B	02-03-1992
		MX 9202630 A	30-06-1992
		NO 843836 A,B,	27-03-1985
		NZ 209494 A	06-03-1987
		US 4582916 A	15-04-1986
		ZA 8407550 A	28-05-1986